

ForPatients

by Roche

Juvenile Idiopathic Arthritis

A Study of Subcutaneously Administered Tocilizumab in Participants With Systemic Juvenile Idiopathic Arthritis

Trial Status
Completed

Trial Runs In
12 Countries

Trial Identifier
NCT01904292 2012-003490-26
WA28118

The source of the below information is the publicly available website [ClinicalTrials.gov](https://clinicaltrials.gov). It has been summarised and edited into simpler language.

Trial Summary:

This open-label, multicenter study will evaluate the pharmacokinetics, pharmacodynamics, and safety of subcutaneously administered tocilizumab in participants with Systemic Juvenile Idiopathic Arthritis (sJIA). Participants with body weight less than (<) 30 kilograms (kg) will receive subcutaneous (SC) tocilizumab dose every 2 weeks (Q2W) and participants with body weight greater than or equal to (\geq) 30 kg will receive weekly (QW), for 52 weeks. Tocilizumab was administered every 10 days until pre-planned interim analysis was performed and changed to Q2W in participants with body weight <30 kg.

Hoffmann-La Roche
Sponsor

Phase 1
Phase

NCT01904292 2012-003490-26 WA28118
Trial Identifiers

Eligibility Criteria:

Gender
All

Age
 ≥ 1 Year & ≤ 17 Years

Healthy Volunteers
No
