

Breast Cancer

A Study of Trastuzumab Emtansine (Kadcyla) Plus Pertuzumab (Perjeta) Following Anthracyclines in Comparison With Trastuzumab (Herceptin) Plus Pertuzumab and a Taxane Following Anthracyclines as Adjuvant Therapy in Participants With Operable HER2-Positive Primary Breast Cancer

Trial Status
Completed

Trial Runs In
36 Countries

Trial Identifier
NCT01966471 2012-004902-82
BO28407

The source of the below information is the publicly available website ClinicalTrials.gov. It has been summarised and edited into simpler language.

Trial Summary:

This two-arm, randomized, open-label, multicenter study will evaluate the efficacy and safety of trastuzumab emtansine in combination with pertuzumab versus trastuzumab in combination with pertuzumab and a taxane as adjuvant therapy in participants with human epidermal growth (HER) factor 2 (HER2)-positive primary invasive breast cancer. Following surgery and anthracycline-based chemotherapy, participants will receive either trastuzumab emtansine at a dose of 3.6 milligrams per kilogram (mg/kg) and pertuzumab at a dose of 420 milligrams (mg) intravenously (IV) every 3 weeks (q3w) or trastuzumab at a dose of 6 mg/kg and pertuzumab at a dose of 420 mg IV q3w in combination with a taxane.

Hoffmann-La Roche
Sponsor

Phase 3
Phase

NCT01966471 2012-004902-82 BO28407
Trial Identifiers

Eligibility Criteria:

Gender
All

Age
≥18 Years

Healthy Volunteers
No
