

Rheumatoid Arthritis

An Efficacy and Safety Study of Subcutaneous Tocilizumab in Combination With Methotrexate (MTX) and as Monotherapy Versus MTX in Participants With Moderate to Severe Rheumatoid Arthritis With Inadequate Response to Current Disease-Modifying Antirheumatic Drug (DMARD) Therapy

Trial Status
Completed

Trial Runs In
1 Countries

Trial Identifier
NCT03155347 YA29359

The source of the below information is the publicly available website ClinicalTrials.gov. It has been summarised and edited into simpler language.

Trial Summary:

This is a randomized, double-blind, multi-center, parallel-group study to evaluate the efficacy and safety of subcutaneous (SC) tocilizumab (162 milligrams [mg] every 2 weeks [Q2W]) given as monotherapy and in combination with MTX versus MTX given as monotherapy, in participants with moderate to severe active rheumatoid arthritis (RA) who have inadequate response to current DMARD therapy. The study comprises a 24-week double-blind treatment phase, followed by a 24-week extension phase.

Hoffmann-La Roche
Sponsor

Phase 3
Phase

NCT03155347 YA29359
Trial Identifiers

Eligibility Criteria:

Gender
All

Age
>= 18 Years & <= 70 Years

Healthy Volunteers
No
