

# ForPatients

by Roche

COVID-19 Pneumonia COVID-19

## Study to Evaluate the Effects of AT-527 in Non-Hospitalized Adult Patients With Mild or Moderate COVID-19

**Trial Status**  
Completed

**Trial Runs In**  
6 Countries

**Trial Identifier**  
NCT04709835 WV43042

---

The source of the below information is the publicly available website [ClinicalTrials.gov](https://clinicaltrials.gov). It has been summarised and edited into simpler language.

### ***Trial Summary:***

This randomized study evaluates the antiviral activity, safety, efficacy and pharmacokinetics of AT-527 versus a placebo in participants with mild or moderate coronavirus disease (COVID-19) who are not hospitalized.

**Hoffmann-La Roche**  
Sponsor

**Phase 2**  
Phase

---

**NCT04709835 WV43042**  
Trial Identifiers

---

### ***Eligibility Criteria:***

**Gender**  
All

**Age**  
≥18 Years

**Healthy Volunteers**  
No

---

**How does the MOONSONG (WV43042) clinical trial work?** This clinical trial is recruiting people who have tested positive for the viral infection COVID-19. In order to take part, patients must have mild or moderate COVID-19 symptoms but must not have severe illness that needs to be treated in hospital.

The purpose of this clinical trial is to compare the effects, good or bad, of AT-527 against placebo in patients with mild or moderate COVID-19. If you take part in this clinical trial, you will receive either AT-527 or a placebo.

**How do I take part in this clinical trial?** To be able to take part in this clinical trial, you must be 18–64 years old and have been diagnosed with COVID-19 at the time of screening. You must also have mild or moderate COVID-19 symptoms that started no more than 5 days before the first dose of study treatment.

# ForPatients

*by Roche*

You must not have COVID-19 symptoms that need to be treated in hospital, have a condition that may increase your risk for developing severe COVID-19 or have previously received treatment for COVID-19. Current or former heavy smokers will also not be able to take part in this trial. If you have a history of some other conditions, you may not be able to take part in this clinical trial.

If you think this clinical trial may be suitable for you and would like to take part, please talk to your doctor. If your doctor thinks that you might be able to take part in this clinical trial, he/she may refer you to the closest clinical trial doctor. They will give you all the information you need to make your decision about taking part in the clinical trial. You can also find the clinical trial locations on this page.

You will have a further COVID-19 test to confirm you definitely have COVID-19 before you take part in the study. During the study you will have some further tests to monitor your health and to understand the effects of the study treatment on your body.

Before starting the clinical trial, you will be told about any risks and benefits of taking part in the trial. You will also be told what other options are available so that you may decide if you still want to take part.

While taking part in the clinical trial, both men and women (if you are not currently pregnant but can become pregnant) will need to either not have heterosexual intercourse or use an effective contraception method, as the effect of the study treatment on a developing baby is currently unknown.

**What treatment will I be given if I join this clinical trial?** Everyone who joins this clinical trial will be enrolled into one of five different groups (A–E), based on when they join the study. In each group, everyone will be split into two treatment arms randomly (like flipping a coin) and given either:

- AT-527 as a pill for five days
- OR placebo as a pill for five days

If you are enrolled into Group A, you will have a 1 in 2 chance of being placed in the AT-527 treatment arm or the placebo treatment arm. If you are enrolled into Groups B, C, D or E, you will have a 3 in 4 chance of being placed in the AT-527 treatment arm and a 1 in 4 chance of being placed in the placebo treatment arm.

- Group A will be given AT-527 or placebo twice per day for five days
- The dose for Groups B, C, D or E will be decided after results from Group A have been reviewed

This is a 'placebo-controlled' clinical trial, which means that one of the groups will be given medicine with no active ingredients (also known as a 'placebo'). Comparing the effects

# ForPatients

*by Roche*

of a study treatment with a placebo helps to show that the study treatment is effective for treating a disease.

Neither you nor your clinical trial doctor can choose or know the treatment arm (AT-527 or placebo) you are in. However, your clinical trial doctor can find out which arm you are in if your safety is at risk.

**How often will I be seen in follow-up appointments and for how long?** You will be given the clinical trial treatment AT-527 or placebo for 5 days.

- If you are in Group A, you will be treated as an outpatient and can take the treatment at home
- If you are in Group B, C, D or E, you will be treated in a clinic and will stay there as a resident for 14 days

You are free to stop this treatment at any time. While being given treatment, you will still be seen regularly by a member of the clinical trial team.

- If you are in Group A, you will be seen every other day at home for the first 7 days so that the clinical trial team can check how you are responding and collect nasal swabs and blood samples
- If you are in Group B, C, D or E, you will be closely monitored throughout the 14 days of your stay so that the clinical trial team can make detailed observations about how your body responds to the treatment

All patients will be asked to complete a diary of their symptoms throughout the study. All patients will receive a follow-up phone call roughly 33 days after the first dose of study treatment, so that the clinical trial team can collect final information.

## What does the MOONSONG (WV43042) clinical trial look like?

### 1. Can I take part in this clinical trial?

If you wish to take part in this clinical trial, you must still meet your usual health and safety requirements for clinical trial participation.



You have not yet had COVID-19 and the laboratory is needed for the study. You will receive the study drug and the placebo. All other tests are done at your own expense or insurance.

You must not be pregnant and you will be screened for any of the following conditions throughout the study.

### 2. What treatment will I be given?



### 3. What happens during the clinical trial?



During your treatment, you will still be responsible for your health and safety in the study and for your own safety.

Group 0 will receive 100 mg of study drug for 14 days. Group 1 will receive 100 mg of study drug for 14 days. Group 2 will receive 100 mg of study drug for 14 days. Group 3 will receive 100 mg of study drug for 14 days.

Group 4, 5, 6 and 7 will be randomly assigned throughout the course of the study to either 100 mg of study drug or placebo.

All patients will be asked to complete a survey at their appointment throughout the study.

All patients will receive a follow-up phone call roughly 14 days after the end of their study or the clinical trial ends for safety reasons.

You will receive the study drug in a clinic and you will not be able to receive the study drug at home.

# ForPatients

*by Roche*

**What happens if I am unable to take part in this clinical trial?** If this clinical trial is not suitable for you, you will not be able to take part. Your doctor will suggest other clinical trials that you may be able to take part in or other treatments that you can be given. You will not lose access to any of your regular care.

For more information about this clinical trial see the For Expert tab on the specific ForPatient page or follow this link to ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT04709835?term=WV43042&draw=2&rank=1>

Trial-identifier: NCT04709835