

Relapsed or Refractory Follicular Lymphoma

A clinical trial to compare mosunetuzumab plus lenalidomide with rituximab plus lenalidomide in people with previously treated follicular lymphoma

A Study Evaluating the Efficacy and Safety of Mosunetuzumab in Combination With Lenalidomide in Comparison to Rituximab in Combination With Lenalidomide With a US Extension of Mosunetuzumab in Combination With Lenalidomide in Participants With Follicular Lymphoma

Trial Status
Recruiting

Trial Runs In
15 Countries

Trial Identifier
NCT04712097 2023-505807-21-00
GO42909

The source of the below information is the publicly available website ClinicalTrials.gov. It has been summarised and edited into simpler language.

Trial Summary:

This study will evaluate the efficacy and safety of mosunetuzumab in combination with lenalidomide (M + Len) compared to rituximab in combination with lenalidomide (R + Len) in participants with relapsed or refractory (R/R) follicular lymphoma (FL) who have received at least one line of prior systemic therapy.

Hoffmann-La Roche
Sponsor

Phase 3
Phase

NCT04712097 2023-505807-21-00 GO42909
Trial Identifiers

Eligibility Criteria:

Gender
All

Age
≥18 Years

Healthy Volunteers
No

[Para descargar el folleto en Español aquí](#)

1. Why is the Celestimo clinical trial needed?

Lymphoma is a type of blood cancer that starts in white blood cells, which are an essential part of our immune system. Follicular lymphoma (FL) is a slow-growing (also sometimes

ForPatients

by Roche

called indolent) form of non-Hodgkin lymphoma (NHL) – the most common type of lymphoma. Standard treatments include radiotherapy, chemotherapy and immunotherapy. Immunotherapies help the body to use its own immune system to fight the cancer. However, FL often comes back after treatment (relapses), and it becomes more difficult to treat with each relapse. Treatments can stop working (known as ‘refractory’ FL) and for people who have had two or more previous therapies, treatment options are currently limited. New treatments are needed to slow or prevent FL getting worse and the chance of relapses. Mosunetuzumab is a type of immunotherapy approved by health authorities for treating relapsed or refractory FL in people who have had at least two treatments before. Mosunetuzumab attaches to a marker called CD20 that is on some types of cancer cells. This brings them closer to cancer-killing immune cells. An immunotherapy called rituximab (that also attaches to CD20 on cancer cells) is an approved standard treatment for FL in combination with another drug called lenalidomide. Lenalidomide can stop cancer cells developing and help the immune system attack cancer cells. Lenalidomide in combination with mosunetuzumab has not been approved by health authorities, and may work well against relapsed or refractory FL.

This clinical trial aims to compare the effects, good or bad, of mosunetuzumab plus lenalidomide with rituximab plus lenalidomide in people with previously treated FL.

2. How does the Celestimo clinical trial work?

This clinical trial is recruiting people with relapsed or refractory FL. People can take part if they need treatment for their FL and have cancer cells that test positive for the CD20 marker. People who take part in this clinical trial (participants) will be given the clinical trial treatment mosunetuzumab plus lenalidomide OR rituximab plus lenalidomide for about 1 year unless their cancer worsens. The clinical trial doctor will see them regularly. These hospital visits will include checks to see how the participant responds to the treatment and any side effects they may have. After the last dose, participants will be seen by the clinical trial doctor every 3 months for up to 4 years or for as long as they agree to it. In addition, participants' medical records will be checked for up to 4 years after the last participant has completed treatment, as long as they agree to it. The total time of participation in the clinical trial will be up to 8 years depending on when they start the trial. Participants can stop trial treatment and leave the clinical trial at any time.

3. What are the main endpoints of the Celestimo clinical trial?

The main clinical trial endpoints (the main results measured to see if the drug has worked) are:

- The length of time between the start of the trial and participants' cancer getting worse (progression-free survival)
- The number of participants whose cancer shrinks or disappears on scans (objective response rate)

ForPatients

by Roche

The other clinical trial endpoints include:

- The number of participants whose cancer disappears on scans (complete response rate) and the length of time this lasts if cancer then gets worse (duration of complete response)
- How long participants live (overall survival)
- The length of time between cancer getting better, then getting worse (duration of response)
- The length of time between the start of trial treatment and cancer symptoms, levels of tiredness and ability to do daily physical tasks getting worse or a new treatment for FL being given
- The number and seriousness of any side effects
- How the body breaks down and gets rid of the trial treatment and its effects on the immune system

4. Who can take part in this clinical trial?

People can take part in this trial if they are at least 18 years old and have had systemic treatment (a treatment that travels through the bloodstream) for FL before. People may not be able to take part in this trial if they have certain other types of cancer or medical conditions. These include autoimmune, heart, liver and lung diseases, certain infections, and being pregnant or breastfeeding. People who have had lenalidomide or certain other treatments before may not be able to take part.

5. What treatment will participants be given in this clinical trial?

Treatment will be given in cycles – a treatment cycle is the treatment and recovery time before the next dose is given. Everyone who joins this clinical trial will be split into two groups randomly (like flipping a coin) with an equal chance of being placed in either group and given either:

- **Mosunetuzumab plus lenalidomide**
 - Mosunetuzumab as an infusion into the vein on Days 1, 8 and 15 for the first 21 days (Cycle 1), and then on Day 1 every 28 days (Cycles 2–12)
 - Lenalidomide as a pill (to be swallowed) once a day for the first 21 days of every 28-day cycle (11 cycles in total), starting from Day 1 of Cycle 2
- **OR rituximab plus lenalidomide**
 - Rituximab as an infusion into the vein on Days 1, 8, 15 and 22 for the first 28 days (Cycle 1), and then on Day 1 of Cycles 3, 5, 7, 9 and 11
 - Lenalidomide as a pill (to be swallowed) once a day for the first 21 days of every 28-day cycle (12 cycles in total), starting from Day 1 of Cycle 1

Everyone who joins this clinical trial extension in the United States will be given **mosunetuzumab plus lenalidomide** in the same way as described above.

ForPatients

by Roche

If a participant experiences a potential side effect called 'cytokine release syndrome' (CRS), they may receive another drug called **tocilizumab** given as an infusion into the vein. This is an open-label trial, which means everyone involved, including the participant and the clinical trial doctor, will know the clinical trial treatment the participant has been given.

6. Are there any risks or benefits in taking part in this clinical trial?

The safety or effectiveness of the experimental treatment or use may not be fully known at the time of the trial. Most trials involve some risks to the participant. However, it may not be greater than the risks related to routine medical care or the natural progression of the health condition. People who would like to participate will be told about any risks and benefits of taking part in the clinical trial, as well as any additional procedures, tests, or assessments they will be asked to undergo. All of these will be described in an informed consent document (a document that provides people with the information they need to decide to volunteer for the clinical trial).

Risks associated with the clinical trial drugs

Participants may have side effects (an unwanted effect of a drug or medical treatment) from the drugs used in this clinical trial. Side effects can be mild to severe, even life-threatening, and vary from person to person. Participants will be closely monitored during the clinical trial; safety assessments will be performed regularly. Participants will be told about the known side effects of mosunetuzumab, lenalidomide, rituximab and tocilizumab and possible side effects based on human and laboratory studies or knowledge of similar drugs. Participants will be told about any known side effects of infusions into the vein (intravenous infusions) and swallowing pills.

Potential benefits associated with the clinical trial

Participants' health may or may not improve from participation in the clinical trial. Still, the information collected may help other people with similar medical conditions in the future.

For more information about this clinical trial see the **For Expert** tab on the specific ForPatient page or follow this link to ClinicalTrials.gov

Trial-identifier: NCT04712097