

Progressive Multiple Sclerosis (PMS)

A Study to Evaluate the Safety, Pharmacokinetics, Pharmacodynamics, and Efficacy of RO7268489 as Add-on Therapy to Ocrelizumab, in Participants With Progressive Forms of Multiple Sclerosis (MS)

Trial Status Not yet recruiting	Trial Runs In	Trial Identifier NCT07282574 2025-521636-10-00 BP46016
---	----------------------	--

The information is taken directly from public registry websites such as ClinicalTrials.gov, EuClinicalTrials.eu, ISRCTN.com, etc., and has not been edited.

Official Title:

A Multi-center, Double-blind, Placebo-controlled, Phase II Study Evaluating the Safety, Pharmacokinetics, Pharmacodynamics, and Efficacy of RO7268489, a Monoacylglycerol Lipase Inhibitor, as Add-on Therapy to Ocrelizumab, in Participants With Progressive Forms of Multiple Sclerosis

Trial Summary:

The main purpose of this study is to assess the efficacy of RO7268489 in adults with progressive multiple sclerosis (PMS) receiving ocrelizumab. After the end of the double-blind period, an open-label (OL) extension may allow eligible participants to receive open-label RO7268489.

Hoffmann-La Roche Sponsor	Phase 2 Phase
-------------------------------------	-------------------------

NCT07282574 2025-521636-10-00 BP46016
Trial Identifiers

Eligibility Criteria:

Gender All	Age #18 Years & # 60 Years	Healthy Volunteers No
----------------------	--------------------------------------	---------------------------------

Inclusion Criteria:

- PMS, in accordance with the revised 2017 McDonald criteria

ForPatients

by Roche

- Expanded disability status scale (EDSS) at screening between 3.0 and 6.0 inclusive

Exclusion Criteria:

- MS relapse during the 6 months preceding the randomization date
- Lack of peripheral venous access
- History of alcohol or other drug abuse, in the opinion of the investigator, within 5 years prior to screening
- Inability to complete an magnetic resonance imaging (MRI)
- Contraindications to ocrelizumab mandatory pre-medications
- Treatment with intravenous immunoglobulin (IV Ig) or plasmapheresis within 12 weeks prior to screening